

# Blood Counts

## CBC

CPT: 85004, 85007, 85008, 85013, 85014, 85018, 85025, 85027, 85032, 85048, 85049

## CMS National Coverage Policy

### Coverage Indications, Limitations, and/or Medical Necessity

Blood counts are used to evaluate and diagnose diseases relating to abnormalities of the blood or bone marrow. These include primary disorders such as anemia, leukemia, polycythemia, thrombocytosis and thrombocytopenia. Many other conditions secondarily affect the blood or bone marrow, including reaction to inflammation and infections, coagulopathies, neoplasms and exposure to toxic substances. Many treatments and therapies affect the blood or bone marrow, and blood counts may be used to monitor treatment effects.

The complete blood count (CBC) includes a hemogram and differential white blood count (WBC). The hemogram includes enumeration of red blood cells, white blood cells, and platelets, as well as the determination of hemoglobin, hematocrit, and indices.

The symptoms of hematological disorders are often nonspecific, and are commonly encountered in patients who may or may not prove to have a disorder of the blood or bone marrow. Furthermore, many medical conditions that are not primarily due to abnormalities of blood or bone marrow may have hematological manifestations that result from the disease or its treatment. As a result, the CBC is one of the most commonly indicated laboratory tests.

In patients with possible hematological abnormalities, it may be necessary to determine the hemoglobin and hematocrit, to calculate the red cell indices, and to measure the concentration of white blood cells and platelets. These measurements are usually performed on a multichannel analyzer that measures all of the parameters on every sample. Therefore, laboratory assessments routinely include these measurements.

### Indications

Indications for a CBC or hemogram include red cell, platelet, and white cell disorders. Examples of these indications are enumerated individually below.

1. Indications for a CBC generally include the evaluation of bone marrow dysfunction as a result of neoplasms, therapeutic agents, exposure to toxic substances, or pregnancy. The CBC is also useful in assessing peripheral destruction of blood cells, suspected bone marrow failure or bone marrow infiltrate, suspected myeloproliferative, myelodysplastic, or lymphoproliferative processes, and immune disorders.
2. Indications for hemogram or CBC related to red cell (RBC) parameters of the hemogram include signs, symptoms, test results, illness, or disease that can be associated with anemia or other red blood cell disorder (e.g., pallor, weakness, fatigue, weight loss, bleeding, acute injury associated with blood loss or suspected blood loss, abnormal menstrual bleeding, hematuria, hematemesis, hematochezia, positive fecal occult blood test, malnutrition, vitamin deficiency, malabsorption, neuropathy, known malignancy, presence of acute or chronic disease that may have associated anemia, coagulation or hemostatic disorders, postural dizziness, syncope, abdominal pain, change in bowel habits, chronic marrow hypoplasia or decreased RBC production, tachycardia, systolic heart murmur, congestive heart failure, dyspnea, angina, nailbed deformities, growth retardation, jaundice, hepatomegaly, splenomegaly, lymphadenopathy, ulcers on the lower extremities).
3. Indications for hemogram or CBC related to red cell (RBC) parameters of the hemogram include signs, symptoms, test results, illness, or disease that can be associated with polycythemia (for example, fever, chills, ruddy skin, conjunctival redness, cough, wheezing, cyanosis, clubbing of the fingers, orthopnea, heart murmur, headache, vague cognitive changes including memory changes, sleep apnea, weakness, pruritus, dizziness, excessive sweating, visual symptoms, weight loss, massive obesity, gastrointestinal bleeding, paresthesias, dyspnea, joint symptoms, epigastric distress, pain and erythema of the fingers or toes, venous or arterial thrombosis, thromboembolism, myocardial infarction, stroke, transient ischemic attacks, congenital heart disease, chronic obstructive pulmonary disease, increased erythropoietin production associated with neoplastic, renal or hepatic disorders, androgen or diuretic use, splenomegaly, hepatomegaly, diastolic hypertension.)

Visit [QuestDiagnostics.com/MLCP](http://QuestDiagnostics.com/MLCP) to view current limited coverage tests, reference guides, and policy information.

To view the complete policy and the full list of codes, please refer to the CMS website reference

[www.cms.gov](http://www.cms.gov) ►

# Blood Counts

CPT: 85004, 85007, 85008, 85013, 85014, 85018, 85025, 85027, 85032, 85048, 85049

---

## CMS National Coverage Policy (continued)

4. Specific indications for CBC with differential count related to the WBC include signs, symptoms, test results, illness, or disease associated with leukemia, infections or inflammatory processes, suspected bone marrow failure or bone marrow infiltrate, suspected myeloproliferative, myelodysplastic or lymphoproliferative disorder, use of drugs that may cause leukopenia, and immune disorders (e.g., fever, chills, sweats, shock, fatigue, malaise, tachycardia, tachypnea, heart murmur, seizures, alterations of consciousness, meningismus, pain such as headache, abdominal pain, arthralgia, odynophagia, or dysuria, redness or swelling of skin, soft tissue bone, or joint, ulcers of the skin or mucous membranes, gangrene, mucous membrane discharge, bleeding, thrombosis, respiratory failure, pulmonary infiltrate, jaundice, diarrhea, vomiting, hepatomegaly, splenomegaly, lymphadenopathy, opportunistic infection, such as oral candidiasis.)
5. Specific indications for CBC related to the platelet count include signs, symptoms, test results, illness, or disease associated with increased or decreased platelet production and destruction, or platelet dysfunction (e.g., gastrointestinal bleeding, genitourinary tract bleeding, bilateral epistaxis, thrombosis, ecchymosis, purpura, jaundice, petechiae, fever, heparin therapy, suspected DIC, shock, pre-eclampsia, neonate with maternal ITP, massive transfusion, recent platelet transfusion, cardiopulmonary bypass, hemolytic uremic syndrome, renal diseases, lymphadenopathy, hepatomegaly, splenomegaly, hypersplenism, neurologic abnormalities, viral or other infection, myeloproliferative, myelodysplastic, or lymphoproliferative disorder, thrombosis, exposure to toxic agents, excessive alcohol ingestion, autoimmune disorder (SLE, RA).
6. Indications for hemogram or CBC related to red cell (RBC) parameters of the hemogram include, in addition to those already listed, thalassemia, suspected hemoglobinopathy, lead poisoning, arsenic poisoning, and spherocytosis.
7. Specific indications for CBC with differential count related to the WBC include, in addition to those already listed, storage diseases; mucopolysaccharidoses, and use of drugs that cause leukocytosis such as G-CSF or CM-CSF.
8. Specific indications for CBC related to platelet count include, in addition to those already listed, May-Hegglin syndrome and Wiskott-Aldrich syndrome.

## Limitations

1. Testing of patients who are asymptomatic, or who do not have a condition that could be expected to result in a hematological abnormality, is screening and is not a covered service.
2. In some circumstances it may be appropriate to perform only a hemoglobin or hematocrit to assess the oxygen carrying capacity of the blood. When the ordering provider requests only a hemoglobin or hematocrit, the remaining components of the CBC are not covered.
3. When a blood count is performed for an end-stage renal disease (ESRD) patient, and is billed outside the ESRD rate, documentation of the medical necessity for the blood count must be submitted with the claim. 4. In some patients presenting with certain signs, symptoms or diseases, a single CBC may be appropriate. Repeat testing may not be indicated unless abnormal results are found, or unless there is a change in clinical condition. If repeat testing is performed, a more descriptive diagnosis code (e.g., anemia) should be reported to support medical necessity. However, repeat testing may be indicated where results are normal in patients with conditions where there is a continued risk for the development of hematologic abnormality.
4. In some patients presenting with certain signs, symptoms or diseases, a single CBC may be appropriate. Repeat testing may not be indicated unless abnormal results are found, or unless there is a change in clinical condition. If repeat testing is performed, a more descriptive diagnosis code (e.g., anemia) should be reported to support medical necessity. However, repeat testing may be indicated where results are normal in patients with conditions where there is a continued risk for the development of hematologic abnormality.

---

Visit [QuestDiagnostics.com/MLCP](https://www.questdiagnostics.com/MLCP) to view current limited coverage tests, reference guides, and policy information.

To view the complete policy and the full list of codes, please refer to the CMS website reference

[www.cms.gov](https://www.cms.gov) ►

# Blood Counts

CPT: 85004, 85007, 85008, 85013, 85014, 85018, 85025, 85027, 85032, 85048, 85049

The ICD10 codes listed below are the top diagnosis codes currently utilized by ordering physicians for the limited coverage test highlighted above that are also listed as medically supportive under Medicare's limited coverage policy. **If you are ordering this test for diagnostic reasons that are not covered under Medicare policy, an Advance Beneficiary Notice form is required.**

**\*Note—Bolded diagnoses below have the highest utilization**

There is a frequency associated with this test. Please refer to the Limitations or Utilization Guidelines section on previous page(s).

Code	Description
D50.9	Iron deficiency anemia, unspecified
D64.9	Anemia, unspecified
E03.9	Hypothyroidism, unspecified
E11.65	Type 2 diabetes mellitus with hyperglycemia
E11.9	Type 2 diabetes mellitus without complications
E53.8	Deficiency of other specified B group vitamins
<b>E55.9</b>	<b>Vitamin D deficiency, unspecified</b>
E78.00	Pure hypercholesterolemia, unspecified
<b>E78.2</b>	<b>Mixed hyperlipidemia</b>
<b>E78.5</b>	<b>Hyperlipidemia, unspecified</b>
<b>I10</b>	<b>Essential (primary) hypertension</b>
I25.10	Atherosclerotic heart disease of native coronary artery without angina pectoris
N39.0	Urinary tract infection, site not specified
R53.83	Other fatigue
R73.01	Impaired fasting glucose
R73.03	Prediabetes
R73.09	Other abnormal glucose
R73.9	Hyperglycemia, unspecified
R79.89	Other specified abnormal findings of blood chemistry
<b>Z79.899</b>	<b>Other long term (current) drug therapy</b>

Visit [QuestDiagnostics.com/MLCP](http://QuestDiagnostics.com/MLCP) to view current limited coverage tests, reference guides, and policy information.

To view the complete policy and the full list of codes, please refer to the CMS website reference

[www.cms.gov](http://www.cms.gov) ►

Last updated: 10/01/22

**Disclaimer:**

This diagnosis code reference guide is provided as an aid to physicians and office staff in determining when an ABN (Advance Beneficiary Notice) is necessary. Diagnosis codes must be applicable to the patient's symptoms or conditions and must be consistent with documentation in the patient's medical record. Quest Diagnostics does not recommend any diagnosis codes and will only submit diagnosis information provided to us by the ordering physician or his/her designated staff. The CPT codes provided are based on AMA guidelines and are for informational purposes only. CPT coding is the sole responsibility of the billing party. Please direct any questions regarding coding to the payer being billed.

**QuestDiagnostics.com**

Quest, Quest Diagnostics, any associated logos, and all associated Quest Diagnostics registered or unregistered trademarks are the property of Quest Diagnostics. All third-party marks—© and ™—are the property of their respective owners. © 2016 Quest Diagnostics Incorporated. All rights reserved.